

Newton's Method

Your work should be presented in the form of a typed report using clear and properly punctuated English. Where appropriate include full program listings and output. If you choose to work in a group of two, please turn in independently prepared reports.

1. Consider Newton's method for solving $f(x) = 0$ where $f(x) = x^2 - 2$ using the starting point $x_0 = 1$.
 - (i) Let $e_n = x_n - \sqrt{2}$ and create a table with three columns showing n , x_n and e_n for $n = 0, 1, \dots, 8$.
 - (ii) A sign of quadratic convergence is that the number of significant digits double at each iteration. Does that happen in this case?
 - (iii) Comment on how rounding error effects the numerical convergence of Newton's method.
 - (iv) Write $|e_{n+1}| = M_n |e_n|^2$ and compute M_n for $n = 1, 2, 3$, and 4. In this case is M_n bigger or less than 1?
 - (v) [Extra Credit and for Math/CS 666] Use multi-precision arithmetic with at least 10 000 digits precision to determine the asymptotic value of M_n when n is large. Can you also find this value analytically?

2. Consider the secant method for solving $f(x) = 0$ where $f(x) = x^2 - 2$ using the starting points $x_0 = 0$ and $x_1 = 1$.
 - (i) Let $e_n = x_n - \sqrt{2}$ and create a table with three columns showing n , x_n and e_n for $n = 0, 1, \dots, 8$.
 - (ii) A sign of quadratic convergence is that the number of significant digits double at each iteration. Does that happen in this case?
 - (iii) According to Wikipedia https://en.wikipedia.org/wiki/Secant_method the order of convergence of the secant method is

$$\alpha = \frac{1 + \sqrt{5}}{2} \approx 1.618,$$

which is less than quadratic. Write $|e_{n+1}| = M_n |e_n|^\alpha$ and compute M_n for $n = 1, 2, \dots, 7$. In this case is M_n bigger or less than 1?

- (iv) [Extra Credit and for Math/CS 666] Prove that the order of convergence of the secant method is α . If you look the proof up, please cite your references and rewrite the proof in your own words.

3. Consider the fixed point iteration for solving $f(x) = 0$ given by $x_{n+1} = h(x_n)$ where

$$h(x) = x - \frac{f(x)f'(x)}{[f'(x)]^2 - f(x)f''(x)}.$$

- (i) Show that $f(x) = 0$ implies $h(x) = x$. Conversely show that if $h(x) = x$ then either $f(x) = 0$ or $f'(x) = 0$.
- (ii) Compute $h'(x)$ and show that

$$h'(x) = \begin{cases} 0 & \text{when } f(x) = 0 \text{ and } f'(x) \neq 0 \\ 2 & \text{when } f(x) \neq 0, f'(x) = 0 \text{ and } f''(x) \neq 0. \end{cases}$$

Conclude that the fixed points of h for which $f(x) = 0$ are stable, but the fixed points for which $f'(x) = 0$ are not.

- (iii) Use this fixed point iteration with $x_0 = 1$ to solve $f(x) = 0$ where $f(x) = x^2 - 2$. Compare the performance of this method with your results for Newton's method.
- (iv) Suppose $f(x) = (x - \xi)^m q(x)$ where $\lim_{x \rightarrow \xi} q(x) \neq 0$. Let $g(x) = x - f(x)/f'(x)$ as in Newton's method and show that

$$\lim_{x \rightarrow \xi} g'(x) = \frac{m-1}{m} \quad \text{and} \quad \lim_{x \rightarrow \xi} h'(x) = 0.$$

Conclude that even when $f'(x) = 0$ this method, unlike Newton's method, has an accelerating rate of convergence as x_n approaches the solution $x = \xi$ to $f(x) = 0$.

4. The function

$$f(x) = 2 \cos(5x) + 2 \cos(4x) + 6 \cos(3x) + 4 \cos(2x) + 10 \cos(x) + 3$$

has two roots on the interval $[0, 3]$; one root is near 1 and the other near 2.

- (i) Use Newton's method $x_{n+1} = g(x_n)$ with $x_0 = 1$ and also with $x_0 = 2$ to approximate these two roots. Use the fact that the exact roots are $\pi/3$ and $2\pi/3$ to compute the error e_n at each iteration for $n = 0, 1, \dots, 18$.
- (ii) Use the method $x_{n+1} = h(x_n)$ with $x_0 = 1$ and again also with $x_0 = 2$ to approximate these two roots. Again use the fact that the exact roots are $\pi/3$ and $2\pi/3$ to compute the error e_n at each iteration for $n = 0, 1, \dots, 18$.
- (iii) Comment on the rate of convergence and the effects of rounding error in the above two computations.

5. [Extra Credit and Math/CS 666] Define

$$x_{n+1} = x_n - \frac{F_n F'_n}{[F'_n]^2 - F_n F''_n}$$

where

$$\begin{aligned} F_n &= f\left(\frac{x_n + 2x_{n-1} + x_{n-2}}{4}\right), \\ F'_n &= \frac{2}{x_n - x_{n-2}} \left\{ f\left(\frac{x_n + x_{n-1}}{2}\right) - f\left(\frac{x_{n-1} + x_{n-2}}{2}\right) \right\}, \\ F''_n &= \frac{2}{x_n - x_{n-2}} \left(\frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} - \frac{f(x_{n-1}) - f(x_{n-2})}{x_{n-1} - x_{n-2}} \right) \end{aligned}$$

to create a secant-method-like approximation of the method given by h that doesn't involve f' and f'' . Study this method both numerically and analytically. Test this method for the functions $f(x) = x^2 - 2$ and

$$f(x) = 2 \cos(5x) + 2 \cos(4x) + 6 \cos(3x) + 4 \cos(2x) + 10 \cos(x) + 3.$$

How does this method compare to the usual secant method? Can you think of an improvement that works better and still doesn't involve f' or f'' ?